



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOLUTION BY A. M. HARDING, University of Arkansas.

Since each of the curves is symmetrical with respect to both axes we shall consider only those areas which lie on the positive side of the x -axis.

$$A_1 = A_2 = \int_0^{a\sqrt{1/2}} (y_1 - y_2)dx = \frac{1}{a^4} \int_0^{a\sqrt{1/2}} [x^2(a^2 - x^2)^{3/2} - x^4(a^2 - x^2)^{1/2}]dx$$

$$= \frac{1}{a^4} \int_0^{a\sqrt{1/2}} x^2 \sqrt{a^2 - x^2} (a^2 - 2x^2) dx.$$

Let

$$x = a \sin \frac{\theta}{2}.$$

Then

$$A_1 = A_2 = \frac{a^2}{8} \int_0^{\pi/2} \sin^2 \theta \cos \theta d\theta = \frac{a^2}{24}.$$

$$A_3 = A_4 = \int_{a\sqrt{1/2}}^a (y_2 - y_1)dx = \frac{1}{a^4} \int_{a\sqrt{1/2}}^a x^2 \sqrt{a^2 - x^2} (2x^2 - a^2) dx.$$

Let $x^2 = a^2 - u^2$. Then

$$A_3 = A_4 = \frac{1}{a^4} \int_0^{a\sqrt{1/2}} u^2 \sqrt{a^2 - u^2} (a^2 - 2u^2) du = \frac{a^2}{24},$$

as above.

$$\frac{1}{2}A_5 = \int_0^{a\sqrt{1/2}} y_2 dx + \int_{a\sqrt{1/2}}^a y_1 dx = \frac{1}{a^4} \int_0^{a\sqrt{1/2}} x^4 \sqrt{a^2 - x^2} dx + \frac{1}{a^4} \int_{a\sqrt{1/2}}^a x^2 (a^2 - x^2)^{3/2} dx.$$

The last integral reduces to

$$\frac{1}{a^4} \int_0^{a\sqrt{1/2}} u^4 \sqrt{a^2 - u^2} du$$

on setting $x^2 = a^2 - u^2$. Hence,

$$\frac{1}{2}A_5 = \frac{2}{a^4} \int_0^{a\sqrt{1/2}} x^4 \sqrt{a^2 - x^2} dx = \frac{a^2}{8} \left(\frac{\pi}{4} - \frac{1}{3} \right),$$

or

$$A_5 = \frac{a^2}{4} \left(\frac{\pi}{4} - \frac{1}{3} \right).$$

Also solved by A. R. NAUER, H. L. OLSON, and the Proposer.

2712 [June, 1918]. Proposed by WILLIAM HOOVER, Columbus, Ohio.

Given the conic $ax^2 + 2hxy + by^2 - 2x = 0$. Find the locus on which lie the four points of intersection of pairs of tangents to the conic from a pair of points on the x -axis equidistant from the origin.

SOLUTION BY A. H. WILSON, Haverford College.

The pair of tangents to the conic $C \equiv ax^2 + 2hxy + by^2 - 2x = 0$ from the point $(\alpha, 0)$ are represented by the equation $\lambda C + l^2 = 0$, where, $l \equiv (a\alpha - 1)x + h\alpha y - \alpha = 0$, is the polar of $(\alpha, 0)$, and $\lambda = \alpha(2 - a\alpha)$. Similarly, if $m \equiv (a\alpha + 1)x + h\alpha y - \alpha = 0$, $\mu = -\alpha(2 + a\alpha)$, $\mu C + m^2 = 0$, represents the tangents from $(-\alpha, 0)$.

The elimination of α from the equations $\lambda C + l^2 = 0$ and $\mu C + m^2 = 0$ is effected at once by subtraction and gives for the required locus the conic $hxy + by^2 - x = 0$.

As a does not occur in this equation, it is the locus described for any one of the conics of the one-parameter family obtained by holding h and b fixed and allowing a to vary.

Also solved by A. M. HARDING and the Proposer.

